The alternating group

We can now use our new tools to understand an important subgroup of Sn called the alternating group, or An.

First we need to define the "sign" of an element in Sn. Recall that we can think of an element of Sn as a permutation of 123...n.

i.e. $\sigma \in S_n$ corresponds to the permutation $\sigma(i)\sigma(2)...\sigma(n)$. Intuitively, we can see that we can get from one permutation to another by successively switching two elements.

To get from 12345 to 32451: (1346)

Def: A 2-cycle is called a transposition.

We can write every m-cycle as a product of 2-cycles as follows:

$$(a_1 a_2 \dots a_m) = (a_i a_m)(a_i a_{m-i}) \dots (a_i a_2)$$

Since every element is the product of cycles, every element of Sn can be written as the product of transpositions.

Note that there is not a unique way to do this! e.g. (12)(13) = (132) = (23)(12) and $1 = (12)(12) = (12)^2(23)^2$.

However, we can determine the parity (i.e. odd or even) of the # of terms, and this will be fixed for a given permutation:

Def: If $\sigma \in S_n$, the sign of σ is 1 if σ can be written as the product of an even # of transpositions, and -1 if σ is the product of an odd #.

Claim: The sign of o GSn is well-defined. That is, any decomposition of o into transpositions has the same parity (odd or even).

We won't prove this, but see D+T= or any other algebra book for a proof.

Prop: The map $\mathcal{E}: S_n \to \mathcal{E}^{\pm} I_s^2$ where each element is sent to its sign is a homomorphism ($\mathcal{E}^{\pm} I_s^2$ is a group under multiplication).

Pf: If o, c & Sn and they can be written as the product of m and h transpositions, respectively, then or can be written as m+n transpositions.

If one of mond n is odd, the other even, then mon is odd, so $\mathcal{E}(\sigma \tau) = -1 = 1 \cdot -1 = \mathcal{E}(\sigma) \mathcal{E}(\tau)$

Otherwise, m+n is even and $\varepsilon(\sigma c) = 1 = (\pm 1)^2 = \varepsilon(\sigma)\varepsilon(c)$. \Box

Def: The kirnel of E is called the <u>alternating group</u> of degree n, denoted An. Thus, An ≤ Sn and An consists of all the even permutations.

Note that a cycle $(a_1 \dots a_m) = (a_1 a_m)(a_1 a_{m-1}) \dots (a_1 a_2)$ is even iff m is odd.

Ex:
$$A_1 = 1$$
, $A_2 = 1$, $A_3 = \{1, (123), (132)\} \in \mathbb{Z}_3$

Ay has 12 elements, the even elements of Sy. Check: it's not isomorphic to D12.

2

ى

In fact, it's isomorphic to the group of symmetries of a tetrahedron!