

A Fan-type degree condition for the existence of disjoint chorded cycles in a graph

RONALD J. GOULD

*Department of Mathematics, Emory University
Atlanta, GA 30322, U.S.A.
rg@emory.edu*

KAZUhide HIROHATA

*Department of Industrial Engineering, Computer Science
National Institute of Technology, Ibaraki College
Hitachinaka, Ibaraki 312-8508, Japan
hirohata@ibaraki-ct.ac.jp*

Abstract

A chord of a cycle is an edge between two non-consecutive vertices of the cycle, and a chorded cycle is a cycle containing at least one chord. Let k be a positive integer. Finkel (2008) proved if G is a graph of order at least $4k$ with the minimum degree at least $3k$, then G contains k disjoint chorded cycles. In this paper, when the order of a graph is sufficiently large, we improve Finkel's result by considering a Fan-type degree condition.

1 Introduction

We consider only finite undirected simple graphs in this paper. A *chord* of a cycle is an edge between two non-consecutive vertices of the cycle, and a *chorded cycle* is a cycle containing at least one chord. The study of cycles has a long and important history. Building on these ideas, investigation of chorded cycles has seen a great deal of development over the last twenty years. Building on this development, we determine a Fan-type degree condition for the existence of disjoint chorded cycles in a graph.

For a graph G , $d_G(u)$ is the degree of a vertex u in G , $\delta(G)$ is the minimum degree of G , and we define $\sigma_2(G) = \min\{d_G(u) + d_G(v) \mid u, v \in V(G), uv \notin E(G)\}$ and $\sigma_2(G) = \infty$ when G is a complete graph. A set of subgraphs of G is said to be *disjoint* if no two of them have any common vertex in G . The distance between two vertices u and v in G is denoted by $\text{dist}_G(u, v)$.

Let k be a positive integer. Corrádi and Hajnal [1] proved if G is a graph of order at least $3k$ with $\delta(G) \geq 2k$, then G contains k disjoint cycles. Enomoto [2] and Wang

[9] independently extended the result of Corrádi and Hajnal showing, if $|G| \geq 3k$ and $\sigma_2(G) \geq 4k - 1$, then G contains k disjoint cycles. Fan [3] considered a new degree condition to improve known sufficient conditions for a graph to be Hamiltonian, called a “Fan-condition”: $\mu(G) = \min\{\max\{d_G(u), d_G(v)\} \mid u, v \in V(G), \text{dist}_G(u, v) = 2\}$, and $\mu(G) = \infty$ when G is a complete graph. Zhang and Yan improved the result of Corrádi and Hajnal by considering a Fan-type condition:

$$\sigma_1^2(G) = \min\{\max\{d_G(u), d_G(v)\} \mid u, v \in V(G), uv \notin E(G)\},$$

and $\sigma_1^2(G) = \infty$ when G is a complete graph.

Theorem 1.1. (Zhang and Yan [10]) *Let k be a positive integer, and let G be a graph of order $n \geq 3k + 2$. If $\sigma_1^2(G) \geq 2k$, then G contains k disjoint cycles.*

Finkel proved the following theorem on the existence of k disjoint chorded cycles.

Theorem 1.2. (Finkel [4]) *Let k be a positive integer, and let G be a graph of order $n \geq 4k$. If $\delta(G) \geq 3k$, then G contains k disjoint chorded cycles.*

In this paper, when the order of a graph is sufficiently large, we improve Finkel’s result (Theorem 1.2) as Corrádi and Hajnal’s result about $\delta(G)$ was improved by Zhang and Yan.

Theorem 1.3. *Let k be a positive integer, and let G be a graph of order $n \geq 12k - 5$. If $\sigma_1^2(G) \geq 3k$, then G contains k disjoint chorded cycles.*

Remark 1.4. The degree condition of Theorem 1.3 is sharp. Consider a complete bipartite graph $G_1 = K_{3k-1, n-3k+1}$, with sufficiently large $n = |G_1|$. Then $\sigma_1^2(G_1) = 3k - 1$. However, G_1 does not contain k disjoint chorded cycles, since any chorded cycle must contain at least three vertices from each partite set. Thus $\sigma_1^2(G) \geq 3k$ for a graph G is necessary.

Since $\sigma_1^2(G) \geq \delta(G)$ for a graph G , Theorem 1.3 strengthens Theorem 1.2 when the order of G is sufficiently large. Moreover, there exists a graph that satisfies the condition of Theorem 1.3, but does not satisfy the condition of Theorem 1.2, and still contains k disjoint chorded cycles. Consider a graph G_2 obtained by deleting an arbitrary edge from a complete bipartite graph $K_{3k, n-3k}$, with sufficiently large $n = |G_2|$. Then $\delta(G_2) = 3k - 1$ and $\sigma_1^2(G_2) = 3k$. By Theorem 1.3, G_2 contains k disjoint chorded cycles, whereas Theorem 1.2 cannot be applied to G_2 . In fact, G_2 contains k disjoint chorded cycles.

For other related results on disjoint chorded cycles in graphs, we refer the reader to [6, 8].

Let G be a graph, H a subgraph of G , and $X \subseteq V(G)$. For $u \in V(G)$, the set of neighbors of u in G is denoted by $N_G(u)$, and we denote $d_G(u) = |N_G(u)|$, $N_H(u) = N_G(u) \cap V(H)$, and $d_H(u) = |N_H(u)|$. Also we denote $d_H(X) = \sum_{u \in X} d_H(u)$. If $H = G$, then $d_G(X) = d_H(X)$. The subgraph of G induced by X is denoted by $\langle X \rangle_G$ or, more simply, by $\langle X \rangle$ if the graph G under discussion is clear. Let $G - X =$

$\langle V(G) - X \rangle_G$ and $G - H = \langle V(G) - V(H) \rangle_G$. If $X = \{x\}$, then we write $G - x$ for $G - X$. If there is no fear of confusion, then we use the same symbol for a graph and its vertex set. For two disjoint $X, Y \subseteq V(G)$, $E(X, Y)$ denotes the set of edges of G connecting a vertex in X and a vertex in Y . Let Q be a path or a cycle with a given orientation and $x \in V(Q)$. Then x^+ denotes the first successor of x on Q and x^- denotes the first predecessor of x on Q . If $x, y \in V(Q)$, then $Q[x, y]$ denotes the path of Q from x to y (including x and y) in the given orientation. We also write $Q(x, y) = Q[x^+, y]$, $Q[x, y) = Q[x, y^-]$, and $Q(x, y) = Q[x^+, y^-]$. If C is a cycle, say $C = x_1x_2 \cdots x_tx_1$, we assume that an orientation of C is given such that x_2 is the successor of x_1 . For two disjoint graphs G_1 and G_2 , $G_1 \cup G_2$ denotes the *union*. For an integer $\ell \geq 3$, a cycle of length ℓ is called an ℓ -cycle. For $x, y \in V(G)$, we denote by xPy a path connecting x and y on G . For terminology and notation not defined here, see [5].

2 Lemmas

For a positive integer r , let $C_1, \dots, C_r$ be r disjoint chorded cycles in a graph G . We say that $C_1, \dots, C_r$ is *minimal* if G does not contain r disjoint chorded cycles $C'_1, \dots, C'_r$ such that $|\bigcup_{i=1}^r V(C'_i)| < |\bigcup_{i=1}^r V(C_i)|$.

Lemma 2.1. *Let G be a graph of order at least 6, and let C be a 4-cycle in G . If $d_C(x) + d_C(y) \geq 7$ for $x, y \in V(G - C)$, then there exists a vertex $z \in V(C)$ such that $xz \in E(G)$ and $\langle V(C - z) \cup \{y\} \rangle$ contains a chorded 4-cycle.*

Proof. Let $C = z_1z_2z_3z_4z_1$. First, suppose that $d_C(x) = 4$. Then $d_C(y) \geq 3$. Without loss of generality, we may assume that $\{z_1, z_2, z_3\} \subseteq N_C(y)$. Then $xz_4 \in E(G)$, and $\langle V(C - z_4) \cup \{y\} \rangle$ contains a 4-cycle $yz_1z_2z_3y$ with chord yz_2 . Next, suppose that $d_C(y) = 4$. Then $d_C(x) \geq 3$. Without loss of generality, we may assume that $xz_1 \in E(G)$. Then $\langle V(C - z_1) \cup \{y\} \rangle$ contains a 4-cycle $yz_2z_3z_4y$ with chord yz_3 . Thus this lemma holds. $\square$

Lemma 2.2. ([7, Lemma 3.2]) *Let $r \geq 1$ be an integer, and let $L = \{C_1, \dots, C_r\}$ be a minimal set of r disjoint chorded cycles in a graph G . If $|C_i| \geq 7$ for some $1 \leq i \leq r$, then C_i has at most two chords. Furthermore, if C_i has two chords, then these chords must be crossing.*

Lemma 2.3. ([7, Lemma 3.3]) *Let $r \geq 1$ be an integer, and let $L = \{C_1, \dots, C_r\}$ be a minimal set of r disjoint chorded cycles in a graph G . Then $d_{C_i}(x) \leq 4$ for any $1 \leq i \leq r$ and any $x \in V(G - \bigcup_{i=1}^r C_i)$. Furthermore, for some $C \in L$ and some $x \in V(G - \bigcup_{i=1}^r C_i)$, if $d_C(x) = 4$, then $|C| = 4$, and if $d_C(x) = 3$, then $|C| \leq 6$.*

Lemma 2.4. ([7, Lemma 3.6]) *Suppose that there exist at least five edges connecting two disjoint paths P_1 and P_2 with $|P_1 \cup P_2| \geq 7$. Then there exists a chorded cycle in $\langle P_1 \cup P_2 \rangle$ not containing at least one vertex of $\langle P_1 \cup P_2 \rangle$.*

Lemma 2.5. ([7, Lemma 3.7]) *Let P_1, P_2 be two disjoint paths, and let u_1, u_2 ($u_1 \neq u_2$) be in that order on P_1 . Suppose that $d_{P_2}(u_i) \geq 2$ for each $i \in \{1, 2\}$. Then there exists a chorded cycle in $\langle P_1[u_1, u_2] \cup P_2 \rangle$.*

Lemma 2.6. *Let $k \geq 2$ be an integer, and let G be a graph. Suppose that G does not contain k disjoint chorded cycles. Let $\{C_1, \dots, C_{k-1}\}$ be a minimal set of $k-1$ disjoint chorded cycles in G , $L = \bigcup_{i=1}^{k-1} C_i$, $H = G - L$, and $X = \{x_1, x_2, x_3, x_4\} \subseteq V(H)$. Suppose that there exists a path connecting x_1 and x_2 in $H - \{x_3, x_4\}$ and there exists a path connecting x_3 and x_4 in $H - \{x_1, x_2\}$. Then $d_{C_i}(X) \leq 12$ for each $1 \leq i \leq k-1$.*

Proof. Suppose not, then $d_{C_{i_0}}(X) \geq 13$ for some $1 \leq i_0 \leq k-1$. By Lemma 2.3, $d_{C_{i_0}}(x_j) \leq 4$ for each $1 \leq j \leq 4$. Then there exists at least one vertex $x \in X$ with $d_{C_{i_0}}(x) = 4$. By Lemma 2.3, $|C_{i_0}| = 4$. Let $C_{i_0} = v_1v_2v_3v_4v_1$. We show the existence of two disjoint chorded cycles in $\langle H \cup C_{i_0} \rangle$, and then G contains k disjoint chorded cycles, a contradiction. It is sufficient to consider the following three cases.

Case 1. X contains three vertices of degree 4 and a vertex of degree at least 1.

Without loss of generality, we may assume that $d_{C_{i_0}}(x_1) \geq 1$ and $d_{C_{i_0}}(x_j) = 4$ for each $2 \leq j \leq 4$. Let $x_1v_1 \in E(G)$. By the assumption of this lemma, there exists a path connecting x_1 and x_2 in $H - \{x_3, x_4\}$. Then $x_1Px_2v_2v_1x_1$ is a cycle with chord x_2v_1 . Since $d_{C_{i_0}}(x_j) = 4$ for each $j \in \{3, 4\}$, $x_3v_3x_4v_4x_3$ is a cycle with chord v_3v_4 . Thus there exist two disjoint chorded cycles in $\langle H \cup C_{i_0} \rangle$.

Case 2. X contains two vertices of degree 4, a vertex of degree 3, and a vertex of degree at least 2.

Without loss of generality, we may assume that $d_{C_{i_0}}(x_1) \geq 2$. First, suppose that $d_{C_{i_0}}(x_2) = 3$. Then, without loss of generality, we may assume that $v_j \in N_{C_{i_0}}(x_2)$ for each $1 \leq j \leq 3$. Assume that $v_1 \in N_{C_{i_0}}(x_1)$. Then $x_1Px_2v_2v_1x_1$ is a cycle with chord x_2v_1 . Since $d_{C_{i_0}}(x_j) = 4$ for each $j \in \{3, 4\}$, $v_3, v_4 \in N_{C_{i_0}}(x_j)$. Then $x_3v_3x_4v_4x_3$ is a cycle with chord v_3v_4 . Thus there exist two disjoint chorded cycles in $\langle H \cup C_{i_0} \rangle$. Hence $v_1 \notin N_{C_{i_0}}(x_1)$. By symmetry, $v_3 \notin N_{C_{i_0}}(x_1)$. Since $d_{C_{i_0}}(x_1) \geq 2$, $v_2 \in N_{C_{i_0}}(x_1)$. Then $x_1Px_2v_1v_2x_1$ is a cycle with chord x_2v_2 . Since $v_3, v_4 \in N_{C_{i_0}}(x_j)$ for each $j \in \{3, 4\}$, $x_3v_3x_4v_4x_3$ is a cycle with chord v_3v_4 . Thus there exist two disjoint chorded cycles in $\langle H \cup C_{i_0} \rangle$.

Next, suppose that $d_{C_{i_0}}(x_3) = 3$ or $d_{C_{i_0}}(x_4) = 3$. Without loss of generality, we may assume that $d_{C_{i_0}}(x_3) = 3$. Then $d_{C_{i_0}}(x_2) = 4$ and $d_{C_{i_0}}(x_4) = 4$. Without loss of generality, we may assume that $v_j \in N_{C_{i_0}}(x_3)$ for each $1 \leq j \leq 3$. Assume that $v_1 \in N_{C_{i_0}}(x_1)$. Then $x_1Px_2v_2v_1x_1$ is a cycle with chord x_2v_1 . Since $d_{C_{i_0}}(x_4) = 4$, $x_3Px_4v_4v_3x_3$ is a cycle with chord x_4v_3 . Thus there exist two disjoint chorded cycles in $\langle H \cup C_{i_0} \rangle$. Hence $v_1 \notin N_{C_{i_0}}(x_1)$. By symmetry, $v_3 \notin N_{C_{i_0}}(x_1)$. Since $d_{C_{i_0}}(x_1) \geq 2$, $v_2 \in N_{C_{i_0}}(x_1)$, and since $d_{C_{i_0}}(x_2) = 4$, $v_1, v_2 \in N_{C_{i_0}}(x_2)$. Then $x_1Px_2v_1v_2x_1$ is a cycle with chord x_2v_2 . Since $d_{C_{i_0}}(x_4) = 4$, $v_3, v_4 \in N_{C_{i_0}}(x_4)$. Then $x_3Px_4v_4v_3x_3$ is a cycle with chord x_4v_3 . Thus there exist two disjoint chorded cycles in $\langle H \cup C_{i_0} \rangle$.

Case 3. X contains a vertex of degree 4 and three vertices of degree 3.

Without loss of generality, we may assume that $d_{C_{i_0}}(x_1) = 4$. Then $d_{C_{i_0}}(x_j) = 3$ for each $2 \leq j \leq 4$. Since $d_{C_{i_0}}(x_3) = 3$, there exist two consecutive vertices $v_{j_0}, v_{j_0+1} \in N_{C_{i_0}}(x_3)$ for some $1 \leq j_0 \leq 4$, with $v_5 = v_1$. Let $v_1, v_2 \in N_{C_{i_0}}(x_3)$. Since $d_{C_{i_0}}(x_4) = 3$, $v_1 \in N_{C_{i_0}}(x_4)$ or $v_2 \in N_{C_{i_0}}(x_4)$. If $v_1 \in N_{C_{i_0}}(x_4)$, then $x_3Px_4v_1v_2x_3$ is

a cycle with chord x_3v_1 . If $v_2 \in N_{C_{i_0}}(x_4)$, then $x_3Px_4v_2v_1x_3$ is a cycle with chord x_3v_2 . Since $d_{C_{i_0}}(x_1) = 4$, $v_3, v_4 \in N_{C_{i_0}}(x_1)$. Since $d_{C_{i_0}}(x_2) = 3$, $v_3 \in N_{C_{i_0}}(x_2)$ or $v_4 \in N_{C_{i_0}}(x_2)$. If $v_3 \in N_{C_{i_0}}(x_2)$, then $x_1Px_2v_3v_4x_1$ is a cycle with chord x_1v_3 . If $v_4 \in N_{C_{i_0}}(x_2)$, then $x_1Px_2v_4v_3x_1$ is a cycle with chord x_1v_4 . Thus there exist two disjoint chorded cycles in $\langle H \cup C_{i_0} \rangle$. $\square$

3 Proof of Theorem 1.3

Let k, G, n be as defined in Theorem 1.3. Since k is a positive integer and $n \geq 12k - 5$, we have $n \geq 7$. Suppose, to the contrary, that G does not contain k disjoint chorded cycles. If G is a complete graph, then Theorem 1.3 holds. Thus we may assume that G is not a complete graph. Suppose that $k = 1$. Then, since $\sigma_1^2(G) \geq 3k$, we have $\sigma_1^2(G) \geq 3$. Thus there exists a vertex $u_0 \in V(G)$ with $d_G(u_0) \geq 3$. Let $\{u_1, u_2, u_3\} \subseteq N_G(u_0)$. Let $P = v_1v_2 \dots v_p$ be a longest path in G . Then $|P| \geq 3$. Suppose that $|P| = 3$. If $u_1u_2 \in E(G)$, then $u_1u_2u_0u_3$ is a longer path than P , a contradiction. Thus we may assume that $u_1u_2 \notin E(G)$. By the $\sigma_1^2(G)$ condition, $d_G(u_1) \geq 3$ or $d_G(u_2) \geq 3$. Since P is a longest path of order 3 in G , this is a contradiction. Thus $|P| \geq 4$. Assume that $v_1v_p \notin E(G)$. By the $\sigma_1^2(G)$ condition, without loss of generality, we may assume that $d_G(v_1) \geq 3$. Since P is a longest path in G , there exists a chorded cycle containing v_1 in $\langle V(P) \rangle$. Thus $v_1v_p \in E(G)$. Let $C = v_1v_2 \dots v_pv_1$. If $v_1v_{p-1} \in E(G)$, then C is a cycle with chord v_1v_{p-1} in G . Thus we may assume that $v_1v_{p-1} \notin E(G)$. By the $\sigma_1^2(G)$ condition, $d_G(v_1) \geq 3$ or $d_G(v_{p-1}) \geq 3$. Since $v_{p-1}v_{p-2} \dots v_1v_p$ is a longest path in G , we have $d_{G-P}(v_{p-1}) = 0$. Thus C is a cycle with chord in G . This is a contradiction.

Suppose that $k \geq 2$. Let G be an edge-maximal counter-example. Let $xy \notin E(G)$ for some $x, y \in V(G)$, and define $G' = G + xy$, the graph obtained from G by adding the edge xy . Since G' is not a counter-example by the edge-maximality of G , G' contains k disjoint chorded cycles $C_1, \dots, C_k$. Without loss of generality, we may assume that $xy \notin \bigcup_{i=1}^{k-1} E(C_i)$, that is, G contains $k - 1$ disjoint chorded cycles. Let $L = \bigcup_{i=1}^{k-1} C_i$, $H = G - L$, and $P_1 = x_1x_2 \dots x_s$ be a longest path in H . We may assume that H does not contain a chorded cycle, otherwise G contains k disjoint chorded cycles, a contradiction. Since G' contains k disjoint chorded cycles, we have $|H| \geq 4$. Choose $C_1, \dots, C_{k-1}$ such that

- (A1) $|L|$ is as small as possible,
- (A2) subject to (A1), $|P_1|$ is as large as possible,
- (A3) subject to (A1) and (A2), $\sum_{x \in V(L)} d_G(x)$ is as large as possible, and
- (A4) subject to (A1)–(A3), $\sum_{x \in V(P_1)} d_G(x)$ is as small as possible.

Since G' is not a counter-example, we note that $|P_1| \geq 4$.

Claim 3.1. *If $d_G(u) \leq 3k - 1$ and $d_G(v) \leq 3k - 1$ for $u, v \in V(G)$, then $uv \in E(G)$.*

Proof. By the $\sigma_1^2(G)$ condition, this claim holds. $\square$

Claim 3.2. $|H| \geq 7$.

Proof. Suppose that $|H| \leq 6$. Assume that $|C_i| \leq 12$ for each $1 \leq i \leq k-1$. Since $|G| \geq 12k-5$ by the assumption, it follows that $|H| \geq (12k-5) - 12(k-1) = 7$, a contradiction. Thus $|C_{i_0}| \geq 13$ for some $1 \leq i_0 \leq k-1$. Without loss of generality, we may assume that C_1 is a longest cycle in L with chord. Then $|C_1| \geq 13$, and $d_{C_1}(v) \leq 2$ for any $v \in V(H)$ by (A1) and Lemma 2.3. By Lemma 2.2, C_1 has at most two chords, and if C_1 has two chords, then these chords must be crossing. Thus, since $|H| \leq 6$ by our assumption, it follows that $|E(H, C_1)| \leq 12$. Let $|C_1| = t (\geq 13)$, and $X = \{v \in V(C_1) \mid d_G(v) \geq 3k\}$. If $|X| \leq t-4$, then there exists a 4-cycle with chords in $\langle V(C_1) \rangle$ by Claim 3.1. This contradicts (A1). Thus we may assume that $|X| \geq t-3$. Then we take a vertex set $X' \subseteq X$ with $|X'| = t-3$.

Suppose that $k = 2$. Then L contains only one cycle C_1 , and since $t \geq 13$,

$$\begin{aligned} |E(X', H)| &= d_G(X') - d_{C_1}(X') \geq 3k(t-3) - (2(t-3) + 4) \\ &= 3kt - 9k - 2t + 2 = 6t - 18 - 2t + 2 \\ &= 4t - 16 \geq 52 - 16 = 36. \end{aligned}$$

Since $12 \geq |E(C_1, H)| \geq |E(X', H)|$, this is a contradiction.

Next, suppose that $k \geq 3$. Since $|H| \leq 6$ and $d_{C_1}(v) \leq 2$ for any $v \in V(H)$,

$$\begin{aligned} |E(X', L - C_1)| &= d_G(X') - d_{C_1}(X') - d_H(X') \\ &\geq 3k(t-3) - (2(t-3) + 4) - 2 \cdot 6 \\ &= 3t(k-1) + t - 9k - 10, \end{aligned}$$

and since $t \geq 13$,

$$\begin{aligned} 3t(k-1) + t - 9k - 10 &= (3t-9)(k-1) + t - 19 \geq (3t-9)(k-1) - 6 \\ &> (3t-9)(k-1) - (3t-9) \\ &= (3t-9)(k-2). \end{aligned}$$

Since $L - C_1$ contains $k-2$ disjoint chorded cycles, we have $|E(X', C')| > 3t-9$ for some C' in $L - C_1$. Let $h = \max\{d_{C'}(v) \mid v \in X'\}$. Let v^* be a vertex of X' such that $d_{C'}(v^*) = h$. If $h \leq 3$, then $|E(X', C')| \leq 3 \times (t-3) = 3t-9$, a contradiction. Thus we may assume that $h \geq 4$. By the maximality of C_1 , $|C'| \leq |C_1| = t$. It follows that $h = d_{C'}(v^*) \leq |C'| \leq t$. Since $t \geq 13$,

$$\begin{aligned} |E(X' - \{v^*\}, C')| &= |E(X', C')| - d_{C'}(v^*) \geq (3t-8) - t \\ &= 2t-8 \geq 18. \end{aligned} \tag{1}$$

Since $h = d_{C'}(v^*) \geq 4$, let v_1, v_2, v_3, v_4 be four distinct neighbors of v^* in that order on C' . Note that v_1, v_2, v_3, v_4 partition C' into four intervals $C'[v_i, v_{i+1})$ for each $1 \leq i \leq 4$, with $v_5 = v_1$. By (1), there exist at least 18 edges from $C_1 - v^*$ to C' . Thus $C'[v_{i_0}, v_{i_0+1})$ for some $1 \leq i_0 \leq 4$ contains at least 5 of these edges. Without loss of generality, we may assume that $i_0 = 4$, that is, $C'[v_4, v_1)$. Then by Lemma 2.4, $\langle (C_1 - v^*) \cup C'[v_4, v_1) \rangle$ contains a chorded cycle C'_1 not containing at least one

vertex of $\langle (C_1 - v^*) \cup C''[v_4, v_1] \rangle$. Note that $C_1'' = v^*C''[v_1, v_3]v^*$ is a cycle with chord v^*v_2 , and it uses no vertices from $C''[v_4, v_1]$. Thus there exist two disjoint chorded cycles C_1', C_1'' in $\langle C_1 \cup C'' \rangle$ such that $|C_1'| + |C_1''| < |C_1| + |C''|$, contradicting (A1). Hence Claim 3.2 holds. $\square$

Claim 3.3. $|P_1| = |H|$.

Proof. Suppose that $|P_1| < |H|$. Let $P_2 = y_1y_2 \dots y_t$ ($t \geq 1$) be a longest path in $H - P_1$. We first claim that

$$\text{there exists a vertex } y_0 \in V(H - P_1) \text{ with } d_H(y_0) \leq 3. \quad (2)$$

By the maximality of P_2 , $d_{H-(P_1 \cup P_2)}(y_i) = 0$ for each $i \in \{1, t\}$. Suppose that $|P_2| \geq 3$. Since H does not contain a chorded cycle, $d_{P_2}(y_i) \leq 2$ for each $i \in \{1, t\}$. By Lemma 2.5, $d_{P_1}(y_{i_0}) \leq 1$ for some $i_0 \in \{1, t\}$. Thus $d_H(y_{i_0}) \leq 2 + 1 = 3$. Suppose that $|P_2| = 2$. Then $t = 2$ and $d_{P_2}(y_i) = 1$ for each $i \in \{1, 2\}$. By Lemma 2.5, $d_{P_1}(y_{i_0}) \leq 1$ for some $i_0 \in \{1, 2\}$. Thus $d_H(y_{i_0}) \leq 1 + 1 = 2$. Suppose that $|P_2| = 1$. Then $t = 1$, $d_{P_2}(y_1) = 0$, and $d_{P_1}(y_1) \leq 2$. Thus $d_H(y_1) \leq 2$. Hence (2) holds.

Next, we claim that

$$d_G(x_i) \geq 3k \text{ for each } i \in \{1, s\}, \text{ and } d_G(y_0) \leq 3k - 1. \quad (3)$$

Suppose that $d_G(y_0) + d_G(x_{i_0}) \geq 6k$ for some $i_0 \in \{1, s\}$. Without loss of generality, we may assume that $i_0 = 1$. Since $d_H(y_0) \leq 3$ by (2), and $d_H(x_1) \leq 2$,

$$\begin{aligned} d_L(y_0) + d_L(x_1) &= d_G(y_0) + d_G(x_1) - (d_H(y_0) + d_H(x_1)) \\ &\geq 6k - (3 + 2) = 6(k - 1) + 1. \end{aligned}$$

Thus $d_{C_{j_0}}(y_0) + d_{C_{j_0}}(x_1) \geq 7$ for some $1 \leq j_0 \leq k - 1$. Then $d_{C_{j_0}}(y_0) \geq 4$ or $d_{C_{j_0}}(x_1) \geq 4$. By (A1) and Lemma 2.3, we have $|C_{j_0}| = 4$. By Lemma 2.1, there exists a vertex $z \in V(C_{j_0})$ such that $x_1z \in E(G)$ and $\langle V(C_{j_0} - z) \cup \{y_0\} \rangle$ contains a chorded 4-cycle, say C_{j_0}'' . Replacing C_{j_0} in L by C_{j_0}'' , we consider the new H' . Then $zx_1x_2 \dots x_s$ is a longer path in H' than P_1 . This contradicts (A2). Thus $d_G(y_0) + d_G(x_i) \leq 6k - 1$ for each $i \in \{1, s\}$. If $d_G(y_0) \leq 3k - 1$, then since $\sigma_1^2(G) \geq 3k$, $d_G(x_i) \geq 3k$ for each $i \in \{1, s\}$. Thus (3) holds. Hence we may assume that $d_G(y_0) \geq 3k$. Since $d_G(y_0) + d_G(x_i) \leq 6k - 1$ for each $i \in \{1, s\}$, we have $d_G(x_i) \leq 3k - 1$. By Claim 3.1, $x_1x_s \in E(G)$. Note that $|P_1| \geq 4$. Since $x_{s-1}x_{s-2} \dots x_1x_s$ is a longest path in H , we have $d_G(x_{s-1}) \leq 3k - 1$ by the above arguments. By Claim 3.1, $x_1x_{s-1} \in E(G)$. Then $x_1x_2 \dots x_sx_1$ is a cycle with chord x_1x_{s-1} , a contradiction. Thus (3) holds.

Next, we claim that

$$d_G(y) \leq 3k - 1 \text{ for any } y \in V(H - P_1), \text{ and } |H - P_1| \leq 3. \quad (4)$$

Suppose that $d_H(y') \geq 4$ for some $y' \in V(H - P_1)$. Since H does not contain a chorded cycle, we have $d_{P_1}(y') \leq 2$, and $d_{H-P_1}(y') \geq 2$. Thus there exists a path of order at least 3 in $H - P_1$. Since $P_2 (= y_1y_2 \dots y_t)$ is a longest path in $H - P_1$, we have

$|P_2| \geq 3$. Since H does not contain a chorded cycle, $d_{P_2}(y_i) \leq 2$ for each $i \in \{1, t\}$. By Lemma 2.5, $d_{P_1}(y_{i_0}) \leq 1$ for some $i_0 \in \{1, t\}$. Note that $d_{H-(P_1 \cup P_2)}(y_{i_0}) = 0$. Thus $d_H(y_{i_0}) \leq 2 + 1 = 3$. Without loss of generality, we may assume that $i_0 = 1$. If $d_G(y_1) \geq 3k$, then since $d_G(x_1) \geq 3k$ by (3), $d_G(y_1) + d_G(x_1) \geq 6k$. Since $d_H(y_1) \leq 3$ and $d_H(x_1) \leq 2$, by the same arguments as the proof of (3), we have $d_{C_{j_0}}(y_1) + d_{C_{j_0}}(x_1) \geq 7$ for some $1 \leq j_0 \leq k - 1$. By Lemma 2.1, there exists a longer path than P_1 , a contradiction. Thus we may assume that $d_G(y_1) \leq 3k - 1$. Since P_1 is a longest path in H and $|P_2| \geq 3$, we have $x_i y_1 \notin E(G)$ for each $i \in \{2, s - 1\}$, and since $d_G(y_1) \leq 3k - 1$, $d_G(x_i) \geq 3k$. We show that

there exist two distinct vertices $u_1, u_2 \in V(H) - \{x_1, x_s\}$ such that $d_G(u_\ell) \geq 3k$, $d_H(u_\ell) \leq 3$ for each $\ell \in \{1, 2\}$, and there exists a path connecting x_1 and u_1 in $H - \{u_2, x_s\}$ and a path connecting u_2 and x_s in $H - \{x_1, u_1\}$. (5)

Suppose that $d_{H-P_1}(x_2) = 0$ and $d_{H-P_1}(x_{s-1}) = 0$. Then we let $u_1 = x_2$ and $u_2 = x_{s-1}$. Since $d_G(y_1) \leq 3k - 1$, $d_G(u_\ell) \geq 3k$ for each $\ell \in \{1, 2\}$. Since H does not contain a chorded cycle, $d_H(u_\ell) \leq 3$ for each $\ell \in \{1, 2\}$.

Suppose that $d_{H-P_1}(x_2) \geq 1$ and $d_{H-P_1}(x_{s-1}) = 0$ or $d_{H-P_1}(x_2) = 0$ and $d_{H-P_1}(x_{s-1}) \geq 1$. By symmetry, we may assume that $d_{H-P_1}(x_2) \geq 1$ and $d_{H-P_1}(x_{s-1}) = 0$. Then there exists a vertex $u \in V(H - P_1)$ with $x_2 u \in E(G)$, and we let $u_1 = u$ and $u_2 = x_{s-1}$. Since P_1 is a longest path in H and $|P_2| \geq 3$, we have $u_1 \notin V(P_2)$. Since $u_1 y_1 \notin E(G)$, $d_G(u_1) \geq 3k$. Since $u_1 x_2 x_3 \dots x_s$ is a longest path in H , we have $d_H(u_1) \leq 2$. Also, we have $d_G(u_2) \geq 3k$ and $d_H(u_2) \leq 3$.

Suppose that $d_{H-P_1}(x_2) \geq 1$ and $d_{H-P_1}(x_{s-1}) \geq 1$. Then there exist vertices $u', u'' \in V(H - P_1)$ such that $x_2 u' \in E(G)$ and $x_{s-1} u'' \in E(G)$. If $u' \neq u''$, then we let $u_1 = u'$ and $u_2 = u''$. Thus we may assume that $u' = u''$. If $|P_1| = 4$ ($s = 4$), then $x_1 x_2 u' x_3 x_4$ is a longer path than P_1 , a contradiction. Thus we may assume that $|P_1| \geq 5$. Suppose that $d_{H-P_1}(x_3) \geq 1$. Let $x_3 w \in E(G)$ for $w \in V(H - P_1)$. If $w = u'$, then $x_1 x_2 w x_3 x_4 \dots x_s$ is a longer path than P_1 . Thus we may assume that $w \neq u'$. Then $x_1 x_2 u' x_{s-1} x_{s-2} \dots x_3 w$ is a longer path than P_1 , a contradiction. Thus $d_{H-P_1}(x_3) = 0$. Since $x_3 y_1 \notin E(G)$, $d_G(x_3) \geq 3k$. We prove that $d_{P_1}(x_3) = 2$. If $x_3 x_1 \in E(G)$, then $x_1 x_2 u' x_{s-1} x_{s-2} \dots x_3 x_1$ is a cycle with chord $x_2 x_3$, a contradiction. If $x_3 x_{r_0} \in E(G)$ for some $5 \leq r_0 \leq s - 1$, then $x_2 x_3 \dots x_{s-1} u' x_2$ is a cycle with chord $x_3 x_{r_0}$, a contradiction. If $x_3 x_s \in E(G)$, then $x_1 x_2 u' x_{s-1} x_{s-2} \dots x_3 x_s$ is a longer path than P_1 , a contradiction. Thus $d_{P_1}(x_3) = 2$. Since $d_{H-P_1}(x_3) = 0$, $d_H(x_3) = 2$. Then we let $u_1 = u'$ and $u_2 = x_3$.

By the above choices of u_1 and u_2 , we note that there exists a path connecting x_1 and u_1 in $H - \{u_2, x_s\}$ and a path connecting u_2 and x_s in $H - \{x_1, u_1\}$. Therefore, (5) holds.

Let $X = \{x_1, u_1, u_2, x_s\}$. Then, by (5),

$$\begin{aligned} d_L(X) &= d_G(X) - d_H(X) \\ &\geq 4 \cdot 3k - (2 + 3 + 3 + 2) = 12k - 10 \\ &= 12(k - 1) + 2. \end{aligned}$$

Hence $d_{C_{i_0}}(X) \geq 13$ for some $1 \leq i_0 \leq k-1$. Since X satisfies the conditions of Lemma 2.6 by (5), this inequality contradicts Lemma 2.6. Hence $d_H(y) \leq 3$ for any $y \in V(H - P_1)$. Similar to (3), we have $d_G(y) \leq 3k-1$ for any $y \in V(H - P_1)$, and by Claim 3.1, $H - P_1$ is a complete graph. Since H does not contain a chorded cycle, we have $|H - P_1| \leq 3$. Thus (4) holds.

Since $d_H(x_i) \leq 2$ for each $i \in \{1, s\}$, we have

$$\begin{aligned} d_L(x_1) + d_L(x_s) &= d_G(x_1) + d_G(x_s) - (d_H(x_1) + d_H(x_s)) \\ &\geq 2 \cdot 3k - (2 + 2) = 6(k-1) + 2. \end{aligned}$$

Thus $d_{C_{i_0}}(x_1) + d_{C_{i_0}}(x_s) \geq 7$ for some $1 \leq i_0 \leq k-1$. Without loss of generality, we may assume that $i_0 = k-1$. Then $d_{C_{k-1}}(x_1) \geq 4$ or $d_{C_{k-1}}(x_s) \geq 4$. By (A1) and Lemma 2.3, we have $|C_{k-1}| = 4$. Let $C_{k-1} = z_1 z_2 z_3 z_4 z_1$. By symmetry, we may assume that $d_{C_{k-1}}(x_1) = 4$ and $\{z_2, z_3, z_4\} \subseteq N_{C_{k-1}}(x_s)$. Then $x_s z_2 z_3 z_4 x_s$ is a 4-cycle with chord $x_s z_3$, and $z_1 x_1 x_2 \dots x_{s-1}$ is a path of the same length as P_1 . Note that $d_G(x_s) \geq 3k$ by (3). Thus, by (A3), we have

$$d_G(z_1) \geq 3k. \quad (6)$$

Subclaim 3.3.1. *The following statements hold.*

- (i) For some $2 \leq i_0 \leq s-1$, $N_{C_{k-1} \cup P_1}(z_1) \subseteq \{z_2, z_3, z_4, x_1, x_{i_0}, x_s\}$, and $N_{H-P_1}(z_1) = \emptyset$.
- (ii) For each $i \in \{2, s-1\}$, $N_{H-P_1}(x_i) = \emptyset$ and $d_G(x_i) \geq 3k$.
- (iii) Let $H_1 = \langle H \cup C_{k-1} \rangle$. Then $d_{H_1}(x_2) + d_{H_1}(x_{s-1}) \leq 10$.

Proof. If $\{x_{i_0}, x_{j_0}\} \subseteq N_{P_1}(z_1)$ for some $2 \leq i_0 < j_0 \leq s-1$, then $\langle P_1 \cup C_{k-1} \rangle$ contains two disjoint chorded cycles, that is, one is a cycle $x_1 x_2 \dots x_{j_0} z_1 x_1$ with chord $x_{i_0} z_1$ and the other is a cycle $x_s z_2 z_3 z_4 x_s$ with chord $x_s z_3$, a contradiction. Thus $N_{C_{k-1} \cup P_1}(z_1) \subseteq \{z_2, z_3, z_4, x_1, x_{i_0}, x_s\}$ for some $2 \leq i_0 \leq s-1$. Suppose that $N_{H-P_1}(z_1) \neq \emptyset$, and let $z_1 y \in E(G)$ for $y \in V(H - P_1)$. Then $x_s z_2 z_3 z_4 x_s$ is a 4-cycle with chord $x_s z_3$, and $y z_1 x_1 x_2 \dots x_{s-1}$ is a longer path than P_1 . This contradicts (A2). Thus $N_{H-P_1}(z_1) = \emptyset$, and (i) holds.

Suppose that $N_{H-P_1}(x_2) \neq \emptyset$. Let $x_2 y \in E(G)$ for $y \in V(H - P_1)$. Then $P'_1 = y x_2 x_3 \dots x_s$ is a path in H with $|P'_1| = |P_1|$. Recall that $d_G(x_1) \geq 3k$ by (3), and $d_G(y) \leq 3k-1$ by (4). Since $\sum_{x \in V(P'_1)} d_G(x) < \sum_{x \in V(P_1)} d_G(x)$, this contradicts (A4). Thus $N_{H-P_1}(x_2) = \emptyset$. By the $\sigma_1^2(G)$ condition and (4), we have $d_G(x_2) \geq 3k$. By symmetry, $N_{H-P_1}(x_{s-1}) = \emptyset$ and $d_G(x_{s-1}) \geq 3k$. Thus (ii) holds.

Suppose that $d_{H_1}(x_2) + d_{H_1}(x_{s-1}) \geq 11$. Since H does not contain a chorded cycle, we have $d_{P_1}(x_i) \leq 3$ for each $i \in \{2, s-1\}$, and since $N_{H-P_1}(x_i) = \emptyset$ by (ii), we have $d_H(x_i) \leq 3$. Thus

$$\begin{aligned} d_{C_{k-1}}(x_2) + d_{C_{k-1}}(x_{s-1}) &= d_{H_1}(x_2) + d_{H_1}(x_{s-1}) - (d_H(x_2) + d_H(x_{s-1})) \\ &\geq 11 - (3 + 3) = 5. \end{aligned}$$

Since $|C_{k-1}| = 4$, $N_{C_{k-1}}(x_2) \cap N_{C_{k-1}}(x_{s-1}) \neq \emptyset$. Thus $z_{i_0} \in N_{C_{k-1}}(x_2) \cap N_{C_{k-1}}(x_{s-1})$ for some $1 \leq i_0 \leq 4$. If $i_0 = 1$, then $x_1x_2 \dots x_{s-1}z_1x_1$ is a cycle with chord x_2z_1 , and $x_s z_2 z_3 z_4 x_s$ is a cycle with chord $x_s z_3$, a contradiction. If $i_0 = 2$, then $x_2x_3 \dots x_s z_2 x_2$ is a cycle with chord $x_{s-1}z_2$, and $x_1z_3z_4z_1x_1$ is a cycle with chord x_1z_4 , a contradiction. If $i_0 \in \{3, 4\}$, then we get a contradiction similar to the case where $i_0 = 2$. Thus (iii) holds. $\square$

Subclaim 3.3.2. For each $1 \leq i \leq k-2$, $2(d_{C_i}(z_1) + d_{C_i}(x_2) + d_{C_i}(x_{s-1})) + d_{C_i}(x_1) + d_{C_i}(x_s) \leq 24$.

Proof. Suppose not. Then, for some $1 \leq i_0 \leq k-2$,

$$2(d_{C_{i_0}}(z_1) + d_{C_{i_0}}(x_2) + d_{C_{i_0}}(x_{s-1})) + d_{C_{i_0}}(x_1) + d_{C_{i_0}}(x_s) \geq 25. \quad (7)$$

Thus $d_{C_{i_0}}(z_1) + d_{C_{i_0}}(x_2) + d_{C_{i_0}}(x_{s-1}) \geq 10$ or $d_{C_{i_0}}(x_1) + d_{C_{i_0}}(x_s) \geq 7$.

First, suppose that $d_{C_{i_0}}(x_1) + d_{C_{i_0}}(x_s) \geq 7$. Then $d_{C_{i_0}}(x_1) \geq 4$ or $d_{C_{i_0}}(x_s) \geq 4$. By (A1) and Lemma 2.3, we have $|C_{i_0}| = 4$. Let $C_{i_0} = w_1w_2w_3w_4w_1$. Without loss of generality, we may assume that $d_{C_{i_0}}(x_1) = 4$ and $\{w_2, w_3, w_4\} \subseteq N_{C_{i_0}}(x_s)$. We claim that $N_{C_{i_0}}(x_2) \cap N_{C_{i_0}}(x_{s-1}) = \emptyset$. Suppose not. We prove that there exist two disjoint chorded cycles in $\langle P_1 \cup C_{i_0} \rangle$, and then G contains k disjoint chorded cycles, a contradiction. Assume that $w_1 \in N_{C_{i_0}}(x_2) \cap N_{C_{i_0}}(x_{s-1})$. Then $x_1x_2 \dots x_{s-1}w_1x_1$ is a cycle with chord x_2w_1 , and $x_s w_2 w_3 w_4 x_s$ is a cycle with chord $x_s w_3$. Assume that $w_2 \in N_{C_{i_0}}(x_2) \cap N_{C_{i_0}}(x_{s-1})$. Then $x_2x_3 \dots x_s w_2 x_2$ is a cycle with chord $x_{s-1}w_2$, and $x_1w_3w_4w_1x_1$ is a cycle with chord x_1w_4 . If $w_{j_0} \in N_{C_{i_0}}(x_2) \cap N_{C_{i_0}}(x_{s-1})$ for some $j_0 \in \{3, 4\}$, then we can prove the existence of two disjoint chorded cycles in $\langle P_1 \cup C_{i_0} \rangle$, similarly. Thus the claim holds. Then $d_{C_{i_0}}(x_2) + d_{C_{i_0}}(x_{s-1}) \leq 4$. Since $|C_{i_0}| = 4$, we have $d_{C_{i_0}}(z_1) \leq 4$ and $d_{C_{i_0}}(x_j) \leq 4$ for each $j \in \{1, s\}$. Hence,

$$2(d_{C_{i_0}}(z_1) + d_{C_{i_0}}(x_2) + d_{C_{i_0}}(x_{s-1})) + d_{C_{i_0}}(x_1) + d_{C_{i_0}}(x_s) \leq 2(4 + 4) + 4 + 4 = 24.$$

This contradicts (7).

Next, suppose that $d_{C_{i_0}}(z_1) + d_{C_{i_0}}(x_2) + d_{C_{i_0}}(x_{s-1}) \geq 10$. Then $d_{C_{i_0}}(v) \geq 4$ for some $v \in \{z_1, x_2, x_{s-1}\}$. We show that $|C_{i_0}| = 4$. If $v \in \{x_2, x_{s-1}\}$, then by (A1) and Lemma 2.3, we have $|C_{i_0}| = 4$. Thus we may assume that $v = z_1$. Recall that $d_{C_{k-1}}(x_1) = 4$. Then $C'_{k-1} = x_1z_2z_3z_4x_1$ is a 4-cycle with chord x_1z_3 . Replacing C_{k-1} in L by C'_{k-1} , we consider the new L' not containing z_1 . Since $d_{C_{i_0}}(z_1) \geq 4$, we have $|C_{i_0}| = 4$ by (A1) and Lemma 2.3. Let $C_{i_0} = w_1w_2w_3w_4w_1$.

We prove that there exist three disjoint chorded cycles in $\langle V(P_1) \cup V(C_{i_0}) \cup V(C_{k-1}) \rangle$. Then G contains k disjoint chorded cycles, a contradiction. Since $|C_{i_0}| = 4$, $d_{C_{i_0}}(z_1) + d_{C_{i_0}}(x_2) + d_{C_{i_0}}(x_{s-1}) \leq 12$.

CASE (A). $d_{C_{i_0}}(z_1) + d_{C_{i_0}}(x_2) + d_{C_{i_0}}(x_{s-1}) = 12$.

In this case, $d_{C_{i_0}}(v) = 4$ for each $v \in \{z_1, x_2, x_{s-1}\}$, and by (7), we have

$$d_{C_{i_0}}(x_1) + d_{C_{i_0}}(x_s) \geq 1. \quad (8)$$

First, we claim that $d_{C_{i_0}}(x_1) = 0$. Assume that $x_1w_1 \in E(G)$. Then $x_1x_2 \dots x_{s-1}w_1x_1$ is a cycle with chord x_2w_1 , $z_1w_2w_3w_4z_1$ is a cycle with chord z_1w_3 , and $x_sz_2z_3z_4x_s$ is a cycle with chord x_sz_3 . Thus $x_1w_1 \notin E(G)$. Similarly, $x_1w_i \notin E(G)$ for each $2 \leq i \leq 4$. Thus the claim holds. Next, we claim that $d_{C_{i_0}}(x_s) = 0$. Assume that $x_sw_1 \in E(G)$. Then $x_2x_3 \dots x_sw_1x_2$ is a cycle with chord $x_{s-1}w_1$, $x_1z_2z_3z_4x_1$ is a cycle with chord x_1z_3 , and $z_1w_2w_3w_4z_1$ is a cycle with chord z_1w_3 . Thus $x_sw_1 \notin E(G)$. Similarly, $x_sw_i \notin E(G)$ for each $2 \leq i \leq 4$. Thus the claim holds. Since $d_{C_{i_0}}(x_i) = 0$ for each $i \in \{1, s\}$, this contradicts (8).

CASE (B). $d_{C_{i_0}}(z_1) + d_{C_{i_0}}(x_2) + d_{C_{i_0}}(x_{s-1}) = 11$.

In this case, $\{d_{C_{i_0}}(z_1), d_{C_{i_0}}(x_2), d_{C_{i_0}}(x_{s-1})\} = \{4, 4, 3\}$, and by (7), we have

$$d_{C_{i_0}}(x_1) + d_{C_{i_0}}(x_s) \geq 3. \quad (9)$$

Case 1. $d_{C_{i_0}}(z_1) = 3$.

In this case, $d_{C_{i_0}}(x_2) = d_{C_{i_0}}(x_{s-1}) = 4$. Without loss of generality, we may assume $\{w_1, w_2, w_3\} \subseteq N_{C_{i_0}}(z_1)$. We claim that $d_{C_{i_0}}(x_1) = 0$. Assume that $x_1w_1 \in E(G)$. Then $x_1x_2w_1z_1x_1$ is a cycle with chord x_1w_1 , $x_{s-1}w_2w_3w_4x_{s-1}$ is a cycle with chord $x_{s-1}w_3$, and $x_sz_2z_3z_4x_s$ is a cycle with chord x_sz_3 . Thus $x_1w_1 \notin E(G)$. Similarly, $x_1w_i \notin E(G)$ for each $i \in \{2, 3\}$. If $x_1w_4 \in E(G)$, then $x_1x_2 \dots x_{s-1}w_4x_1$ is a cycle with chord x_2w_4 , $z_1w_1w_2w_3z_1$ is a cycle with chord z_1w_2 , and $x_sz_2z_3z_4x_s$ is a cycle with chord x_sz_3 . Thus the claim holds. If $x_sw_4 \in E(G)$, then $x_2x_3 \dots x_sw_4x_2$ is a cycle with chord $x_{s-1}w_4$, $x_1z_2z_3z_4x_1$ is a cycle with chord x_1z_3 , and $z_1w_1w_2w_3z_1$ is a cycle with chord z_1w_2 . Thus $x_sw_4 \notin E(G)$. Since $d_{C_{i_0}}(x_1) = 0$, we have $d_{C_{i_0}}(x_s) \geq 3$ by (9). Then $N_{C_{i_0}}(x_s) = \{w_1, w_2, w_3\}$. Thus $x_2x_3 \dots x_{s-1}w_3w_4x_2$ is a cycle with chord $x_{s-1}w_4$, $x_1z_2z_3z_4x_1$ is a cycle with chord x_1z_3 , and $x_sw_1z_1w_2x_s$ is a cycle with chord w_1w_2 .

Case 2. $d_{C_{i_0}}(x_2) = 3$.

In this case, $d_{C_{i_0}}(z_1) = d_{C_{i_0}}(x_{s-1}) = 4$. Without loss of generality, we may assume $\{w_1, w_2, w_3\} \subseteq N_{C_{i_0}}(x_2)$. We claim that $d_{C_{i_0}}(x_1) = 0$. Assume that $x_1w_1 \in E(G)$. Then $x_1x_2 \dots x_{s-1}w_1x_1$ is a cycle with chord x_2w_1 , $z_1w_2w_3w_4z_1$ is a cycle with chord z_1w_3 , and $x_sz_2z_3z_4x_s$ is a cycle with chord x_sz_3 . Thus $x_1w_1 \notin E(G)$. Similarly, $x_1w_j \notin E(G)$ for each $j \in \{2, 3\}$. If $x_1w_4 \in E(G)$, then $x_1z_1w_3w_4x_1$ is a cycle with chord z_1w_4 , $x_2x_3 \dots x_{s-1}w_2w_1x_2$ is a cycle with chord $x_{s-1}w_1$, and $x_sz_2z_3z_4x_s$ is a cycle with chord x_sz_3 . Thus the claim holds. If $x_sw_1 \in E(G)$, then $x_2x_3 \dots x_sw_1x_2$ is a cycle with chord $x_{s-1}w_1$, $x_1z_2z_3z_4x_1$ is a cycle with chord x_1z_3 , and $z_1w_2w_3w_4z_1$ is a cycle with chord z_1w_3 . Thus $x_sw_1 \notin E(G)$. Similarly, $x_sw_i \notin E(G)$ for each $i \in \{2, 3\}$. Thus $d_{C_{i_0}}(x_s) \leq 1$. Since $d_{C_{i_0}}(x_1) = 0$, $d_{C_{i_0}}(x_s) \geq 3$ by (9). This is a contradiction.

Case 3. $d_{C_{i_0}}(x_{s-1}) = 3$.

We can prove this case similar to Case 2.

CASE (C). $d_{C_{i_0}}(z_1) + d_{C_{i_0}}(x_2) + d_{C_{i_0}}(x_{s-1}) = 10$.

In this case, by (7), we have

$$d_{C_{i_0}}(x_1) + d_{C_{i_0}}(x_s) \geq 5. \quad (10)$$

Case 1. $\{d_{C_{i_0}}(z_1), d_{C_{i_0}}(x_2), d_{C_{i_0}}(x_{s-1})\} = \{4, 4, 2\}$.

Subcase 1.1. $d_{C_{i_0}}(z_1) = 2$.

In this case, $d_{C_{i_0}}(x_2) = d_{C_{i_0}}(x_{s-1}) = 4$.

(i) The vertex z_1 is adjacent to consecutive vertices on C_{i_0} .

In this case, without loss of generality, we may assume that $w_1, w_2 \in N_{C_{i_0}}(z_1)$. Since $x_s z_2 z_3 z_4 x_s$ is a cycle with chord $x_s z_3$, we show the existence of two disjoint chorded cycles in $\langle (V(P_1) - \{x_s\}) \cup V(C_{i_0}) \cup \{z_1\} \rangle$. Assume that $x_1 w_1 \in E(G)$. Then $x_1 z_1 w_2 w_1 x_1$ is a cycle with chord $z_1 w_1$, and $x_2 w_3 x_{s-1} w_4 x_2$ is a cycle with chord $w_3 w_4$. Thus $x_1 w_1 \notin E(G)$. Similarly, $x_1 w_2 \notin E(G)$. Assume that $x_1 w_3 \in E(G)$. Then $x_1 w_3 w_4 x_2 x_1$ is a cycle with chord $x_2 w_3$, and $x_{s-1} w_1 z_1 w_2 x_{s-1}$ is a cycle with chord $w_1 w_2$. Thus $x_1 w_3 \notin E(G)$. Similarly, $x_1 w_4 \notin E(G)$. Thus $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$.

(ii) The vertex z_1 is not adjacent to consecutive vertices on C_{i_0} .

In this case, without loss of generality, we may assume that $w_1, w_3 \in N_{C_{i_0}}(z_1)$. Assume that $x_1 w_1 \in E(G)$. Then $x_1 x_2 w_1 z_1 x_1$ is a cycle with chord $x_1 w_1$, and $x_{s-1} w_2 w_3 w_4 x_{s-1}$ is a cycle with chord $x_{s-1} w_3$. Also, $x_s z_2 z_3 z_4 x_s$ is a cycle with chord $x_s z_3$. Thus $x_1 w_1 \notin E(G)$. Similarly, $x_1 w_3 \notin E(G)$. Then $d_{C_{i_0}}(x_1) \leq 2$. By (10), $d_{C_{i_0}}(x_s) \geq 3$. Suppose that $x_1 w_2 \in E(G)$. Then $x_1 x_2 w_1 w_2 x_1$ is a cycle with chord $x_2 w_2$. Since $d_{C_{i_0}}(x_s) \geq 3$, $x_s w_3 \in E(G)$ or $x_s w_4 \in E(G)$. If $x_s w_3 \in E(G)$, then $x_{s-1} x_s w_3 w_4 x_{s-1}$ is a cycle with chord $x_{s-1} w_3$, and if $x_s w_4 \in E(G)$, then $x_{s-1} x_s w_4 w_3 x_{s-1}$ is a cycle with chord $x_{s-1} w_4$. Also, we consider a chorded cycle C_{k-1} . Thus $x_1 w_2 \notin E(G)$. By symmetry, $x_1 w_4 \notin E(G)$. Thus $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$.

Subcase 1.2. $d_{C_{i_0}}(x_2) = 2$.

In this case, $d_{C_{i_0}}(z_1) = d_{C_{i_0}}(x_{s-1}) = 4$. Since $x_s z_2 z_3 z_4 x_s$ is a cycle with chord $x_s z_3$, we show the existence of two disjoint chorded cycles in $\langle (V(P_1) - \{x_s\}) \cup V(C_{i_0}) \cup \{z_1\} \rangle$.

(i) The vertex x_2 is adjacent to consecutive vertices on C_{i_0} .

In this case, without loss of generality, we may assume that $w_1, w_2 \in N_{C_{i_0}}(x_2)$. Assume that $x_1 w_1 \in E(G)$. Then $x_1 x_2 \dots x_{s-1} w_1 x_1$ is a cycle with chord $x_2 w_1$, and $z_1 w_2 w_3 w_4 z_1$ is a cycle with chord $z_1 w_3$. Thus $x_1 w_1 \notin E(G)$. Similarly, $x_1 w_2 \notin E(G)$. Assume that $x_1 w_3 \in E(G)$. Then $x_1 w_3 w_4 z_1 x_1$ is a cycle with chord $z_1 w_3$, and $x_2 w_1 x_{s-1} w_2 x_2$ is a cycle with chord $w_1 w_2$. Thus $x_1 w_3 \notin E(G)$. Similarly, $x_1 w_4 \notin E(G)$. Thus $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$.

(ii) The vertex x_2 is not adjacent to consecutive vertices on C_{i_0} .

In this case, without loss of generality, we may assume that $w_1, w_3 \in N_{C_{i_0}}(x_2)$. Assume that $x_1 w_1 \in E(G)$. Then $x_1 x_2 \dots x_{s-1} w_1 x_1$ is a cycle with chord $x_2 w_1$, and

$z_1w_2w_3w_4z_1$ is a cycle with chord z_1w_3 . Thus $x_1w_1 \notin E(G)$. Similarly, $x_1w_3 \notin E(G)$. Assume that $x_1w_2 \in E(G)$. Then $x_1w_2w_3z_1x_1$ is a cycle with chord z_1w_2 , and $x_2x_3 \dots x_{s-1}w_4w_1x_2$ is a cycle with chord $x_{s-1}w_1$. Thus $x_1w_2 \notin E(G)$. Similarly, $x_1w_4 \notin E(G)$. Thus $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$.

Subcase 1.3. $d_{C_{i_0}}(x_{s-1}) = 2$.

In this case, we get a contradiction by the same arguments as Subcase 1.2.

Case 2. $\{d_{C_{i_0}}(z_1), d_{C_{i_0}}(x_2), d_{C_{i_0}}(x_{s-1})\} = \{4, 3, 3\}$.

Subcase 2.1. $d_{C_{i_0}}(z_1) = 4$.

In this case, $d_{C_{i_0}}(x_2) = d_{C_{i_0}}(x_{s-1}) = 3$. Without loss of generality, we may assume that $N_{C_{i_0}}(x_2) = \{w_1, w_2, w_3\}$. Since $x_s z_2 z_3 z_4 x_s$ is a cycle with chord $x_s z_3$, we show the existence of two disjoint chorded cycles in $\langle (V(P_1) - \{x_s\}) \cup V(C_{i_0}) \cup \{z_1\} \rangle$.

(i) $N_{C_{i_0}}(x_{s-1}) = \{w_1, w_2, w_3\}$.

Assume that $x_1w_1 \in E(G)$. Then $x_1x_2 \dots x_{s-1}w_1x_1$ is a cycle with chord x_2w_1 , and $z_1w_2w_3w_4z_1$ is a cycle with chord z_1w_3 . Thus $x_1w_1 \notin E(G)$. Similarly, $x_1w_3 \notin E(G)$. Assume that $x_1w_2 \in E(G)$. Then $x_1x_2 \dots x_{s-1}w_2x_1$ is a cycle with chord x_2w_2 , and $z_1w_3w_4w_1z_1$ is a cycle with chord z_1w_4 . Thus $x_1w_2 \notin E(G)$. Assume that $x_1w_4 \in E(G)$. Then $x_1z_1w_3w_4x_1$ is a cycle with chord z_1w_4 , and $x_2w_1x_{s-1}w_2x_2$ is a cycle with chord w_1w_2 . Thus $x_1w_4 \notin E(G)$. Therefore, $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$.

(ii) $N_{C_{i_0}}(x_{s-1}) = \{w_1, w_2, w_4\}$ or $N_{C_{i_0}}(x_{s-1}) = \{w_2, w_3, w_4\}$.

By symmetry, it is sufficient to prove the case where $N_{C_{i_0}}(x_{s-1}) = \{w_1, w_2, w_4\}$. Assume that $x_1w_1 \in E(G)$. Then $x_1x_2 \dots x_{s-1}w_1x_1$ is a cycle with chord x_2w_1 , and $z_1w_2w_3w_4z_1$ is a cycle with chord z_1w_3 . Thus $x_1w_1 \notin E(G)$. Assume that $x_1w_2 \in E(G)$. Then $x_1x_2 \dots x_{s-1}w_2x_1$ is a cycle with chord x_2w_2 , and $z_1w_3w_4w_1z_1$ is a cycle with chord z_1w_4 . Thus $x_1w_2 \notin E(G)$. Assume that $x_1w_3 \in E(G)$. Then $x_1x_2w_2w_3x_1$ is a cycle with chord x_2w_3 , and $z_1w_4x_{s-1}w_1z_1$ is a cycle with chord w_1w_4 . Thus $x_1w_3 \notin E(G)$. Assume that $x_1w_4 \in E(G)$. Then $x_1z_1w_3w_4x_1$ is a cycle with chord z_1w_4 , and $x_2x_3 \dots x_{s-1}w_2w_1x_2$ is a cycle with chord $x_{s-1}w_1$. Thus $x_1w_4 \notin E(G)$. Therefore, $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$.

(iii) $N_{C_{i_0}}(x_{s-1}) = \{w_1, w_3, w_4\}$.

Assume that $x_1w_1 \in E(G)$. Then $x_1x_2w_2w_1x_1$ is a cycle with chord x_2w_1 , and $z_1w_3x_{s-1}w_4z_1$ is a cycle with chord w_3w_4 . Thus $x_1w_1 \notin E(G)$. Similarly, $x_1w_3 \notin E(G)$. Assume that $x_1w_2 \in E(G)$. Then $x_1x_2w_3w_2x_1$ is a cycle with chord x_2w_2 , and $z_1w_1x_{s-1}w_4z_1$ is a cycle with chord w_1w_4 . Thus $x_1w_2 \notin E(G)$. Assume that $x_1w_4 \in E(G)$. Then $x_1z_1w_3w_4x_1$ is a cycle with chord z_1w_4 , and $x_2x_3 \dots x_{s-1}w_1w_2x_2$ is a cycle with chord x_2w_1 . Thus $x_1w_4 \notin E(G)$. Therefore, $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$.

Subcase 2.2. $d_{C_{i_0}}(x_2) = 4$.

In this case, $d_{C_{i_0}}(x_{s-1}) = d_{C_{i_0}}(z_1) = 3$. Without loss of generality, we may assume that $N_{C_{i_0}}(x_{s-1}) = \{w_1, w_2, w_3\}$. Since $x_s z_2 z_3 z_4 x_s$ is a cycle with chord $x_s z_3$, we show the existence of two disjoint chorded cycles in $\langle (V(P_1) - \{x_s\}) \cup V(C_{i_0}) \cup \{z_1\} \rangle$.

(i) $N_{C_{i_0}}(z_1) = \{w_1, w_2, w_3\}$.

Assume that $x_1 w_1 \in E(G)$. Then $x_1 z_1 w_2 w_1 x_1$ is a cycle with chord $z_1 w_1$, and $x_2 x_3 \dots x_{s-1} w_3 w_4 x_2$ is a cycle with chord $x_2 w_3$. Thus $x_1 w_1 \notin E(G)$. Similarly, $x_1 w_3 \notin E(G)$. Assume that $x_1 w_2 \in E(G)$. Then $x_1 z_1 w_1 w_2 x_1$ is a cycle with chord $z_1 w_2$, and $x_2 x_3 \dots x_{s-1} w_3 w_4 x_2$ is a cycle with chord $x_2 w_3$. Thus $x_1 w_2 \notin E(G)$. Assume that $x_1 w_4 \in E(G)$. Then $x_1 x_2 w_3 w_4 x_1$ is a cycle with chord $x_2 w_4$, and $x_{s-1} w_1 z_1 w_2 x_{s-1}$ is a cycle with chord $w_1 w_2$. Thus $x_1 w_4 \notin E(G)$. Therefore, $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$.

(ii) $N_{C_{i_0}}(z_1) = \{w_1, w_2, w_4\}$ or $N_{C_{i_0}}(z_1) = \{w_2, w_3, w_4\}$.

By symmetry, it is sufficient to prove the case where $N_{C_{i_0}}(z_1) = \{w_1, w_2, w_4\}$. Assume that $x_1 w_1 \in E(G)$. Then $x_1 z_1 w_4 w_1 x_1$ is a cycle with chord $z_1 w_1$, and $x_2 x_3 \dots x_{s-1} w_3 w_2 x_2$ is a cycle with chord $x_{s-1} w_2$. Thus $x_1 w_1 \notin E(G)$. Assume that $x_1 w_2 \in E(G)$. Then $x_1 w_2 w_1 z_1 x_1$ is a cycle with chord $z_1 w_2$, and $x_2 x_3 \dots x_{s-1} w_3 w_4 x_2$ is a cycle with chord $x_2 w_3$. Thus $x_1 w_2 \notin E(G)$. Assume that $x_1 w_3 \in E(G)$. Then $x_1 x_2 w_4 w_3 x_1$ is a cycle with chord $x_2 w_3$, and $x_{s-1} w_1 z_1 w_2 x_{s-1}$ is a cycle with chord $w_1 w_2$. Thus $x_1 w_3 \notin E(G)$. Assume that $x_1 w_4 \in E(G)$. Then $x_1 x_2 w_3 w_4 x_1$ is a cycle with chord $x_2 w_4$, and $x_{s-1} w_1 z_1 w_2 x_{s-1}$ is a cycle with chord $w_1 w_2$. Thus $x_1 w_4 \notin E(G)$. Therefore, $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$.

(iii) $N_{C_{i_0}}(z_1) = \{w_1, w_3, w_4\}$.

Assume that $x_1 w_1 \in E(G)$. Then $x_1 z_1 w_4 w_1 x_1$ is a cycle with chord $z_1 w_1$, and $x_2 x_3 \dots x_{s-1} w_2 w_3 x_2$ is a cycle with chord $x_{s-1} w_3$. Thus $x_1 w_1 \notin E(G)$. Similarly, $x_1 w_3 \notin E(G)$. Assume that $x_1 w_2 \in E(G)$. Then $x_1 x_2 \dots x_{s-1} w_2 x_1$ is a cycle with chord $x_2 w_2$, and $z_1 w_3 w_4 w_1 z_1$ is a cycle with chord $z_1 w_4$. Thus $x_1 w_2 \notin E(G)$. Assume that $x_1 w_4 \in E(G)$. Then $x_1 z_1 w_1 w_4 x_1$ is a cycle with chord $z_1 w_4$, and $x_2 x_3 \dots x_{s-1} w_3 w_2 x_2$ is a cycle with chord $x_{s-1} w_2$. Thus $x_1 w_4 \notin E(G)$. Therefore, $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$.

Subcase 2.3. $d_{C_{i_0}}(x_{s-1}) = 4$.

In this case, $d_{C_{i_0}}(x_2) = d_{C_{i_0}}(z_1) = 3$. Without loss of generality, we may assume that $N_{C_{i_0}}(x_2) = \{w_1, w_2, w_3\}$.

(i) $N_{C_{i_0}}(z_1) = \{w_1, w_2, w_3\}$.

Assume that $x_1 w_1 \in E(G)$. Then $x_1 z_1 w_2 w_1 x_1$ is a cycle with chord $z_1 w_1$, and $x_2 x_3 \dots x_{s-1} w_4 w_3 x_2$ is a cycle with chord $x_{s-1} w_3$. Also, $x_s z_2 z_3 z_4 x_s$ is a cycle with chord $x_s z_3$. Thus $x_1 w_1 \notin E(G)$. Similarly, $x_1 w_3 \notin E(G)$. Assume that $x_1 w_2 \in E(G)$. Then $x_1 z_1 w_1 w_2 x_1$ is a cycle with chord $z_1 w_2$, and $x_2 x_3 \dots x_{s-1} w_4 w_3 x_2$ is a cycle with chord $x_{s-1} w_3$. Also, $x_s z_2 z_3 z_4 x_s$ is a cycle with chord $x_s z_3$. Thus $x_1 w_2 \notin E(G)$. Therefore, $d_{C_{i_0}}(x_1) \leq 1$. By (10), $d_{C_{i_0}}(x_s) \geq 4$. Then $x_1 z_2 z_3 z_4 x_1$ is a cycle with

chord x_1z_3 , $x_2w_1z_1w_2x_2$ is a cycle with chord w_1w_2 , and $x_{s-1}x_sw_3w_4x_{s-1}$ is a cycle with chord $x_{s-1}w_3$, a contradiction.

(ii) $N_{C_{i_0}}(z_1) = \{w_1, w_2, w_4\}$ or $N_{C_{i_0}}(z_1) = \{w_2, w_3, w_4\}$.

By symmetry, it is sufficient to prove the case where $N_{C_{i_0}}(z_1) = \{w_1, w_2, w_4\}$. Since $x_sz_2z_3z_4x_s$ is a cycle with chord x_sz_3 , we show the existence of two disjoint chorded cycles in $\langle(V(P_1) - \{x_s\}) \cup V(C_{i_0}) \cup \{z_1\}\rangle$. Assume that $x_1w_1 \in E(G)$. Then $x_1z_1w_2w_1x_1$ is a cycle with chord z_1w_1 , and $x_2x_3 \dots x_{s-1}w_4w_3x_2$ is a cycle with chord $x_{s-1}w_3$. Thus $x_1w_1 \notin E(G)$. Assume that $x_1w_2 \in E(G)$. Then $x_1x_2w_3w_2x_1$ is a cycle with chord x_2w_2 , and $z_1w_1x_{s-1}w_4z_1$ is a cycle with chord w_1w_4 . Thus $x_1w_2 \notin E(G)$. Assume that $x_1w_3 \in E(G)$. Then $x_1x_2w_2w_3x_1$ is a cycle with chord x_2w_3 , and $z_1w_1x_{s-1}w_4z_1$ is a cycle with chord w_1w_4 . Thus $x_1w_3 \notin E(G)$. Assume that $x_1w_4 \in E(G)$. Then $x_1z_1w_1w_4x_1$ is a cycle with chord z_1w_4 , and $x_2w_2x_{s-1}w_3x_2$ is a cycle with chord w_2w_3 . Thus $x_1w_4 \notin E(G)$. Therefore, $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$.

(iii) $N_{C_{i_0}}(z_1) = \{w_1, w_3, w_4\}$.

Since $x_sz_2z_3z_4x_s$ is a cycle with chord x_sz_3 , we show the existence of two disjoint chorded cycles in $\langle(V(P_1) - \{x_s\}) \cup V(C_{i_0}) \cup \{z_1\}\rangle$. Assume that $x_1w_1 \in E(G)$. Then $x_1x_2w_1z_1x_1$ is a cycle with chord x_1w_1 , and $x_{s-1}w_2w_3w_4x_{s-1}$ is a cycle with chord $x_{s-1}w_3$. Thus $x_1w_1 \notin E(G)$. Similarly, $x_1w_3 \notin E(G)$. Assume that $x_1w_2 \in E(G)$. Then $x_1x_2w_1w_2x_1$ is a cycle with chord x_2w_2 , and $z_1w_3x_{s-1}w_4z_1$ is a cycle with chord w_3w_4 . Thus $x_1w_2 \notin E(G)$. Assume that $x_1w_4 \in E(G)$. Then $x_1z_1w_1w_4x_1$ is a cycle with chord z_1w_4 , and $x_2w_2x_{s-1}w_3x_2$ is a cycle with chord w_2w_3 . Thus $x_1w_4 \notin E(G)$. Therefore, $d_{C_{i_0}}(x_1) = 0$. By (10), $d_{C_{i_0}}(x_s) \geq 5$. This contradicts $|C_{i_0}| = 4$. □

Let

$$\begin{aligned} H_1 &= \langle H \cup C_{k-1} \rangle, \\ d_1 &= \sum_{i=1}^{k-2} \{2(d_{C_i}(z_1) + d_{C_i}(x_2) + d_{C_i}(x_{s-1})) + d_{C_i}(x_1) + d_{C_i}(x_s)\}, \text{ and} \\ d_2 &= 2(d_{H_1}(z_1) + d_{H_1}(x_2) + d_{H_1}(x_{s-1})) + d_{H_1}(x_1) + d_{H_1}(x_s). \end{aligned}$$

By (3), (6), and Subclaim 3.3.1 (ii), we have $d_G(z_1) \geq 3k$ and $d_G(x_j) \geq 3k$ for each $j \in \{1, 2, s-1, s\}$. By Subclaim 3.3.2, $d_1 \leq 24(k-2)$. Since $|C_{k-1}| = 4$, $d_{H_1}(x_1) = d_H(x_1) + d_{C_{k-1}}(x_1) \leq 2 + 4 = 6$. Similarly, $d_{H_1}(x_s) \leq 6$. By Subclaims 3.3.1 (i) and (iii), $d_2 \leq 2(6 + 10) + 6 + 6 = 44$. Thus we have

$$\begin{aligned} 24k &\leq 2(d_G(z_1) + d_G(x_2) + d_G(x_{s-1})) + d_G(x_1) + d_G(x_s) \\ &= d_1 + d_2 \leq 24(k-2) + 44 = 24k - 4. \end{aligned}$$

This is a contradiction. ■

Claim 3.4. *Let $X = \{x \in V(P_1) \mid d_G(x) \geq 3k\}$. Then $|X| \leq 3$.*

Proof. Suppose that $|X| \geq 4$. We claim that we may assume that $d_G(x_1) \geq 3k$. Suppose that $x_1x_s \notin E(G)$. By the $\sigma_1^2(G)$ condition, without loss of generality, we may assume that $d_G(x_1) \geq 3k$. Suppose that $x_1x_s \in E(G)$. By Claims 3.2 and 3.3, we have $|P_1| \geq 7$. Let $C' = x_1x_2 \dots x_sx_1$. Then $|C'| \geq 7$. Since H does not contain a chorded cycle, C' does not have a chord. Thus $x_1x_{s-1} \notin E(G)$. By the $\sigma_1^2(G)$ condition, without loss of generality, we may assume that $d_G(x_1) \geq 3k$. Thus the claim holds. Since $x_1 \in X$, there exist at least three vertices x_p, x_q, x_r ($2 \leq p < q < r \leq s$) in X . We choose x_r such that r is as large as possible. If $r \leq s - 4$, then $\langle \{x_{s-3}, x_{s-2}, x_{s-1}, x_s\} \rangle$ is a complete subgraph of order 4 in H by Claim 3.1, and contains a chorded cycle, a contradiction. Thus we may assume that $r \geq s - 3$.

Case 1. $r \in \{s - 3, s - 1, s\}$.

We prove that $d_{P_1}(x_r) \leq 3$ for each $r \in \{s - 3, s - 1, s\}$. Suppose that $r \in \{s - 1, s\}$. Since H does not contain a chorded cycle, we have $d_{P_1}(x_r) \leq 3$. Suppose that $r = s - 3$. Since $d_G(x_{r'}) \leq 3k - 1$ for each $s - 2 \leq r' \leq s$, we have $x_{s-2}x_s \in E(G)$ by Claim 3.1. If $x_{s-3}x_{s-1} \in E(G)$, then $x_{s-3}x_{s-2}x_sx_{s-1}x_{s-3}$ is a cycle with chord $x_{s-2}x_{s-1}$. If $x_{s-3}x_s \in E(G)$, then $x_{s-3}x_{s-2}x_{s-1}x_sx_{s-3}$ is a cycle with chord $x_{s-2}x_s$. Since H does not contain a chorded cycle, we have $d_{P_1}(x_{s-3}) \leq 3$. Thus $d_{P_1}(x_r) \leq 3$ for each $r \in \{s - 3, s - 1, s\}$, and by Claim 3.3, $d_H(x_r) = d_{P_1}(x_r) \leq 3$. Then

$$\begin{aligned} d_L(x_1) + d_L(x_r) &= d_G(x_1) + d_G(x_r) - (d_H(x_1) + d_H(x_r)) \\ &\geq 3k + 3k - (2 + 3) = 6(k - 1) + 1. \end{aligned}$$

Thus $d_{C_{i_0}}(x_1) + d_{C_{i_0}}(x_r) \geq 7$ for some $1 \leq i_0 \leq k - 1$. Without loss of generality, we may assume that $i_0 = k - 1$. Then $d_{C_{k-1}}(x_1) \geq 4$ or $d_{C_{k-1}}(x_r) \geq 4$. By (A1) and Lemma 2.3, we have $|C_{k-1}| = 4$. Let $C_{k-1} = z_1z_2z_3z_4z_1$. Suppose that $d_{C_{k-1}}(x_1) \geq 4$ (in fact, $d_{C_{k-1}}(x_1) = 4$) and $\{z_2, z_3, z_4\} \subseteq N_{C_{k-1}}(x_r)$. If $d_{C_{k-1}}(x_r) \geq 4$, then we can prove this case by the same arguments as $d_{C_{k-1}}(x_1) \geq 4$.

We prove that $d_G(z_1) \geq 3k$. Suppose that $d_G(z_1) < 3k$. Assume that $r \in \{s - 3, s - 1\}$. Then $C'_{k-1} = x_1z_2z_3z_4x_1$ is a 4-cycle with chord x_1z_3 , and we consider the new L' containing C'_{k-1} instead of C_{k-1} . Let $H' = G - L'$. By the maximality of r , we have $d_G(x_s) < 3k$. By Claim 3.1, $x_sx_1 \in E(G)$. Thus $x_2x_3 \dots x_sx_1$ is a longest path (the same length as P_1) in H' . Since $d_G(x_1) \geq 3k$ and $d_G(z_1) < 3k$, this contradicts (A3). Assume that $r = s$. Then $C''_{k-1} = x_sx_2z_3z_4x_s$ is a 4-cycle with chord x_sx_3 , and we consider the new L'' containing C''_{k-1} instead of C_{k-1} . Let $H'' = G - L''$. Then $z_1x_1x_2 \dots x_{s-1}$ is a longest path (the same length as P_1) in H'' . Since $d_G(x_s) \geq 3k$ and $d_G(z_1) < 3k$, this contradicts (A3). Thus $d_G(z_1) \geq 3k$. Let $H_1 = \langle H \cup C_{k-1} \rangle$. We prove that $d_{H_1}(z_1) \leq 6$. Suppose that $d_{P_1}(z_1) \geq 4$. Let $x_{i_1}, x_{i_2}, x_{i_3} \in N_{P_1-x_1}(z_1)$ for $i_1 < i_2 < i_3$. Then $z_1x_{i_1}x_{i_1+1} \dots x_{i_3}z_1$ is a cycle with chord $z_1x_{i_2}$, and $x_1z_2z_3z_4x_1$ is a cycle with chord x_1z_3 , a contradiction. Thus $d_{P_1}(z_1) \leq 3$. Since $d_{C_{k-1}}(z_1) \leq 3$, $d_{H_1}(z_1) \leq 6$.

We have two observations as follows. Since $d_{P_1}(x_i) \leq 3$ for each $i \in \{2, s - 1\}$, we have

$$d_{P_1}(x_2) + d_{P_1}(x_{s-1}) \leq 6. \quad (11)$$

We prove that for each $1 \leq j \leq s-1$,

$$d_{P_1}(x_j) + d_{P_1}(x_{j+1}) \leq 7. \quad (12)$$

Suppose that $d_{P_1}(x_{j_0}) + d_{P_1}(x_{j_0+1}) \geq 8$ for some $1 \leq j_0 \leq s-1$. Then $d_{P_1}(x_{j_0}) = 4$ and $d_{P_1}(x_{j_0+1}) = 4$. Let $x_{j_0}x_{t_2}, x_{j_0+1}x_{t_1} \in E(G)$ for $j_0 + 1 < t_2$ and $t_1 < j_0$. Then $x_{t_1}x_{t_1+1} \dots x_{j_0}x_{t_2}x_{t_2-1} \dots x_{j_0+1}x_{t_1}$ is a cycle with chord $x_{j_0}x_{j_0+1}$, a contradiction. Thus (12) holds. We show that

$$x_p \text{ and } x_q \text{ can be chosen so that } d_{P_1}(x_p) + d_{P_1}(x_q) \leq 7. \quad (13)$$

Let $Y = \{y \in V(P_1) \mid d_G(y) \leq 3k-1\}$. By Claim 3.1, $|Y| \leq 3$. Suppose that $r = s$. If $|Y| = 0$, then we let $x_p = x_2$ and $x_q = x_{s-1}$ by (11). If $|Y| \in \{1, 2\}$, then since $|P_1| \geq 7$, we can choose the desired vertices x_p, x_q by (11) and (12). Suppose that $|Y| = 3$. Then Y consists of three consecutive vertices on P_1 , otherwise, there exists a chorded cycle in H by Claim 3.1. Since $|P_1| \geq 7$, we can choose the desired vertices x_p, x_q by (11) and (12). Thus (13) holds.

Suppose that $r = s-1$. Then $x_s \in Y$. Let $Y' = Y - \{x_s\}$. If $|Y'| \in \{0, 1\}$, then since $|P_1| \geq 7$, we can choose the desired vertices x_p, x_q by (12). If $|Y'| = 2$, then since $r = s-1$, there exists a chorded cycle in H by Claim 3.1. Suppose that $r = s-3$. Then $x_j \in Y$ for each $s-2 \leq j \leq s$. Since $|P_1| \geq 7$, we can choose the desired vertices x_p, x_q by (12).

Since H_1 does not contain two disjoint chorded cycles, $N_{C_{k-1}}(x_p) \cap N_{C_{k-1}}(x_q) = \emptyset$ similar to the proof of Subclaim 3.3.1 (iii). Thus $d_{C_{k-1}}(x_p) + d_{C_{k-1}}(x_q) \leq 4$. By (13), $d_{H_1}(x_p) + d_{H_1}(x_q) \leq 7 + 4 = 11$. By regarding x_p, x_q, x_r as x_2, x_{s-1}, x_s , we have $2(d_{C_i}(z_1) + d_{C_i}(x_p) + d_{C_i}(x_q)) + d_{C_i}(x_1) + d_{C_i}(x_r) \leq 24$ for each $1 \leq i \leq k-2$ similar to Subclaim 3.3.2. Also, $d_{H_1}(x_1) = d_{P_1}(x_1) + d_{C_{k-1}}(x_1) \leq 2 + 4 = 6$ and $d_{H_1}(x_r) = d_{P_1}(x_r) + d_{C_{k-1}}(x_r) \leq 3 + 4 = 7$. Thus we have $2(d_{H_1}(z_1) + d_{H_1}(x_p) + d_{H_1}(x_q)) + d_{H_1}(x_1) + d_{H_1}(x_r) \leq 2(6 + 11) + 6 + 7 = 47$. Thus $2(d_G(z_1) + d_G(x_p) + d_G(x_q)) + d_G(x_1) + d_G(x_r) \leq 24(k-2) + 47 = 24k-1$. Since $d_G(z_1) \geq 3k$ and $d_G(x_j) \geq 3k$ for each $j \in \{1, p, q, r\}$, this is a contradiction.

Case 2. $r = s-2$.

If $d_G(x_{i_0}) \leq 3k-1$ for some $1 \leq i_0 \leq s-3$, then $x_{i_0}x_{s-1}, x_{i_0}x_s \in E(G)$ by Claim 3.1, and $x_{i_0}x_{i_0+1} \dots x_sx_{i_0}$ is a cycle with chord $x_{i_0}x_{s-1}$, a contradiction. Thus we may assume that $d_G(x_i) \geq 3k$ for each $1 \leq i \leq s-3$. If $d_{P_1}(x_{s-2}) \leq 3$, then we get a contradiction by the same arguments as Case 1. Thus we may assume that $d_{P_1}(x_{s-2}) = 4$. Then $x_{s-2}x_s \in E(G)$. If $d_{P_1}(x_{s-3}) = 4$, then $x_{s-3}x_{s-1} \in E(G)$ or $x_{s-3}x_s \in E(G)$. In both cases, $\langle \{x_{s-3}, x_{s-2}, x_{s-1}, x_s\} \rangle$ contains a chorded cycle. Thus we may assume that $d_{P_1}(x_{s-3}) \leq 3$. Then we get a contradiction by the same arguments as Case 1. □

Let $Y = \{y \in V(P_1) \mid d_G(y) \leq 3k-1\}$. By Claims 3.2, 3.3, and 3.4, we have $|Y| \geq 4$. By Claim 3.1, $\langle Y \rangle$ is a complete subgraph in H . Thus H contains a chorded cycle, a contradiction. ■

Acknowledgements

The authors would like to thank the referees for their careful reading and helpful suggestions. The first author is supported by the Heilbrun Distinguished Emeritus Fellowship from Emory University. The second author is supported by JSPS KAKENHI Grant Number JP23K03207.

References

- [1] K. Corrádi and A. Hajnal, On the maximal number of independent circuits in a graph, *Acta Math. Acad. Sci. Hungar.* 14 (1963), 423–439.
- [2] H. Enomoto, On the existence of disjoint cycles in a graph, *Combinatorica* 18 (4) (1998), 487–492.
- [3] G.H. Fan, New sufficient conditions for cycles in graphs, *J. Combin. Theory Ser. B* 37 (3) (1984), 221–227.
- [4] D. Finkel, On the number of independent chorded cycles in a graph, *Discrete Math.* 308 (22) (2008), 5265–5268.
- [5] R. J. Gould, “Graph Theory”, Dover Pub. Inc. Mineola, N.Y. 2012.
- [6] R. J. Gould, Results and problems on chorded cycles: a survey, *Graphs Combin.* 38 (6) (2022), 189.
- [7] R. J. Gould, K. Hirohata and A.K. Rorabaugh, On independent triples and vertex-disjoint chorded cycles in graphs, *Australas. J. Combin.* 77(3) (2020), 355–372.
- [8] M. Santana and M. Van Bonn, Sharp minimum degree conditions for disjoint doubly chorded cycles, *J. Comb.* 15 (2) (2024), 217–282.
- [9] H. Wang, On the maximum number of independent cycles in a graph, *Discrete Math.* 205 (1-3) (1999), 183–190.
- [10] J. Zhang and J. Yan, Disjoint cycles and 2-factors with Fan-type condition in a graph, *Discrete Math.* 345 (4) (2022), 112789.

(Received 1 Sep 2025; revised 21 Mar 2026)